



Principles and guidelines in the management of ankle fractures in adults

Journal:	<i>Journal of Perioperative Practice</i>
Manuscript ID	PPJ-20-0029.R5
Manuscript Type:	Review
Keywords:	Ankle fracture, Perioperative, Classification, Complications, Trauma and orthopaedic surgery
Abstract:	Ankle fractures are common injuries that have many physical and psychosocial complications. As a result, it is important to be aware of how these patients present and are managed perioperatively. Detailed guidelines from NICE and the British Orthopaedic Association have been produced on this topic, including recent developments such as the decision to weight-bear early after surgery, and the use of virtual fracture clinics. This article provides an overview of the key perioperative factors that need to be considered in cases of ankle fracture and the relevant clinical guidelines.

SCHOLARONE™
Manuscripts

Principles and guidelines in the management of ankle fractures in adults

Abstract

Ankle fractures are common injuries that have many physical and psychosocial complications. As a result, it is important to be aware of how these patients present and are managed perioperatively. Detailed guidelines from NICE and the British Orthopaedic Association have been produced on this topic, including recent developments such as the decision to weight-bear early after surgery, and the use of virtual fracture clinics. This article provides an overview of the key perioperative factors that need to be considered in cases of ankle fracture and the relevant clinical guidelines.

Key words

Ankle fracture; perioperative; complications; orthopaedics

Key phrases

1. The ankle is a complex region and many different injuries can occur
2. Ankle fractures are associated with significant morbidity and occasionally mortality
3. It is important to be aware of the physical and psychosocial complications of ankles fractures
4. Early weight-bearing may improve rehabilitation in stable ankle fractures

Introduction

Ankle fractures are common, representing 14% of all fractures requiring hospitalisation (Jennison & Brinsden 2019). Between 2004-2005 and 2013-2014, there were 332,617 hospital admissions in England due to ankle fractures, accounting for 10% of hospital bed stays (Jennison & Brinsden 2019). Ankle fractures affect the lateral malleolus in 55% of cases and commonly occur due to sports injuries in adolescents (22%), or low-energy falls in later years (61%) (Elsoe et al 2018). They are associated with significant morbidity in all age groups and have a one-year mortality rate of 11.9% after hospitalisation in patients over the age of 65 years (Hsu et al 2015).


The National Institute for Health and Care Excellence (NICE) has published detailed guidelines on how to assess, monitor and manage ankle fractures (NICE 2016a; NICE 2016b). Partly based on these guidelines, the British Orthopaedic Association (BOA) has


published standards for practice in the management of ankle fractures, known as the British Orthopaedic Association Standards for Trauma and Orthopaedics (BOAST) guidelines (BOA 2016). This article provides an overview of ankle fractures and their relevant clinical guidelines, and presents them in a readable format that follows the patient journey. The aim of this article is to raise awareness of these injuries, particularly in the perioperative period, to optimise the care that these patients receive.

Clinical Anatomy

The ankle joint is a synovial mortise and tenon made up of the articular surface of the tibia, both malleoli, and the talus (Moore et al 2013). It works with the subtalar joint to act as a modified hinge which can plantar-flex, dorsi-flex, glide and roll (McKeon & Hoch 2019). To add stability, the ankle is also bound by three lateral ligaments and a strong medial deltoid ligament (Moore et al 2013). This is therefore a complex region with many potential sites of injury (Figure 1).

Figure 1: Anatomy of the ankle





Pre-operative considerations

Presentation, history and examination

Ankle fracture patients typically present with immediate-onset ankle pain that increases with activity, localised or generalised swelling, bruising, joint deformity and an inability to weight-bear (Slimmon & Brukner 2010). Severe ankle sprains may present with similar features. Therefore, history-taking and examination are particularly important in distinguishing between different types of ankle injury.

In the history, it is important to ascertain the timing of injury, mechanism (inversion, eversion, plantar flexion or dorsiflexion), severity (force and velocity involved) and whether the patient was able to bear weight immediately after. The presence and onset of ankle swelling also need to be established, as acute-onset swelling may suggest bleeding (Slimmon & Brukner 2010). A past medical history should include previous ankle injuries which may have predisposed the joint to fracture (Slimmon & Brukner 2010). In addition, co-morbidities that might influence treatment choice and outcome should be documented, e.g. pre-existing mobility impairment, diabetes mellitus, peripheral neuropathy, peripheral vascular disease, osteoporosis, renal disease, smoking and alcohol overuse (BOA 2016).

Examination should include inspection for deformity, bruising, effusion and open wounds (Lampridis et al 2018; BOA & BAPRAS 2017). Palpation is performed in a methodical sequence across both the lateral and medial ankle, as well as proximal fibula (for a potential

Maisonneuve injury) (Lampridis et al 2018). Tenderness is suggestive of an underlying fracture, rather than sprain. A neurovascular assessment of the foot should also be performed, compared with the contralateral limb, and the results documented (NICE 2016b). This includes sensation over the dorsal and plantar surfaces of the foot, distal pulses and capillary refill in all digits (Mordecai et al 2011).

Initial management

The latest NICE guidelines emphasise the importance of regularly assessing pain in ankle fracture patients, using a scale suitable for the patient's age, developmental stage and cognitive function (NICE 2016a). Oral paracetamol should be offered for mild pain, oral paracetamol and codeine for moderate pain, and intravenous (IV) paracetamol with IV morphine titrated to effect for severe pain (NICE 2016a). IV opioids should be used with caution in elderly patients and non-steroidal anti-inflammatory drugs should be avoided (NICE 2016a).

Clinically deformed ankles require urgent reduction and splinting (BOA 2016). Radiographs should not be performed before reduction if they will cause an unacceptable delay (BOA 2016). Reduction minimises the risk of skin necrosis and reduces pain and swelling (Mordecai et al 2011). After reduction, the neurovascular status should be reassessed and documented (BOA 2016). Any fracture should be stabilised in a well-fitted backslab cast or splint, with the limb elevated and a post-reduction X-ray arranged to confirm adequate alignment (Mordecai et al 2011; BOA 2016).

Patients with open fractures, which make up 1.5% of all ankle fractures, should have any gross contaminants removed and their injury photographed (Bugler et al 2015; BOA & BAPRAS 2017). The fracture site should then be covered with a saline-soaked sterile dressing and wrapped loosely with an occlusive film whilst awaiting debridement surgery (BOA & BAPRAS 2017). Intravenous antibiotic prophylaxis should be given as soon as possible, preferably within 1 hour of injury (BOA & BAPRAS 2017). Debridement should occur immediately for highly contaminated open fractures, within 12 hours for high-energy open fractures that are not highly contaminated, and within 24 hours for all other open fractures (BOA & BAPRAS 2017). Fixation and definitive soft tissue cover should be performed at the same time where possible, or if not possible, within 72 hours of injury (NICE 2016b).

1
2
3 106
4
5 107 Following initial management, the ankle should be immobilised using a splint. Then,
6
7 108 definitive treatment is chosen depending on the stability of the joint (Lampridis et al 2018).
8
9 109 Patients with stable fractures or co-morbidities rendering them unfit for surgery are treated
10
11 110 conservatively with analgesia and immobilisation using a splint, short-leg cast or walker boot
12
13 111 for a minimum of six weeks (Moredecai et al 2011; Lampridis et al 2018). In addition to this,
14
15 112 the BOA recommends close contact casts for patients over 60 years of age as an alternative to
16
17 113 surgery if reduction can be maintained by the cast (BOA 2016). These patients exhibit poor
18
19 114 bone and soft tissue quality which leads to poor surgical outcomes (Srinivasan & Moran
20
21 115 2001). By comparison, unstable ankle fractures are generally treated surgically as discussed
22
23 116 later. This may be preceded by temporary external fixation to allow soft tissue swelling to
24
25 117 settle down in some cases, e.g. high-energy fractures of the distal tibial articular surface
26
27 118 (known as pilon fractures) (Shah et al 2019).
28
29 119


30
31 120 *Imaging*
32
33 121 It can often be difficult to clinically distinguish between bony and ligamentous ankle injuries.
34
35 122 Historically, this meant that most patients would receive costly, time-consuming X-rays and
36
37 123 unnecessary radiation exposure (Stiell et al 1992). As a result, the ankle Ottawa rules were
38
39 124 created to guide which patients needed to be imaged. These rules state that an X-ray is only
40
41 125 required if there is malleolar pain and any one of the following: an inability to weight bear
42
43 126 immediately and in the emergency department for four steps; bony tenderness of the posterior
44
45 127 tibia/fibula; or bony tenderness at the tips of the medial/lateral malleolus (Stiell et al 1992).
46
47 128 The literature suggests that the Ottawa ankle rules have a sensitivity of almost 100%, with a
48
49 129 specificity of 39.8%, and have reduced unnecessary radiographs by 30-40% (Bachmann et al
50
51 130 2003).
52
53 131

54
55 132 Imaging should include lateral and mortise view radiographs (Vangsness et al 1994). A
56
57 133 mortise view is an anteroposterior image of the leg in 15-20 degrees of internal rotation. If
58
59 134 radiographs are equivocal or significant ligament/tendon injury is expected, a magnetic
60
135 resonance imaging scan is the imaging of choice (Sawant & Sanghvi 2018). Computed
136
137 tomography (CT) scanning is useful for the assessment of complex ankle fractures, especially
138
139 those involving the posterior malleolus and/or comminuted malleolar fractures (Lampridis et
al 2018; Leung et al 2016).

140 Classification

141 Ankle fractures can be classified using the Danis-Weber classification system (shown in
 142 Figure 2). Type A fractures involve the lateral malleolus distal to the tibiofibular
 143 syndesmosis. They are usually stable and are managed conservatively (Donken et al 2012).
 144 Type B fractures occur at the level of the syndesmosis and have variable stability. They can
 145 be treated both conservatively and surgically (Donken et al 2012). Type C fractures occur
 146 proximal to the syndesmosis, which is often disrupted, and are unstable. In these cases, there
 147 is usually a concurrent fracture of the medial malleolus or injury to the deltoid ligament.
 148 Type C fractures are likely to require open reduction and internal fixation (ORIF) (Donken et
 149 al 2012).

Figure 2: The Danis-Weber classification system



Right foot viewed posteriorly. (A) Type A. (B) Type B. (C) Type C with disrupted syndesmosis.

151
 152 Other less commonly used ankle fracture classifications include the Lauge-Hansen
 153 classification which focuses on the mechanism of injury, as determined by a series of

cadaver-based experiments (Lauge-Hansen 1954). The classification of ankle fractures is a dynamic topic and new schemes have been proposed as recently as 2019 (Briet et al 2019).

Operative considerations

The decision to operate

Unstable fractures are treated surgically in patients deemed fit enough to undergo surgery. These generally include: fractures with syndesmotic disruption (Weber type C and some type B fractures); displaced fractures; unstable bi-/tri-malleolar fractures; and fractures with joint incongruity or talar subluxation (Mordecai et al 2011). The decision to treat posterior malleolar fractures is more controversial, with some authors recommending surgical fixation for fracture fragments that range from 10% to 25% of the distal tibial articular surface (Odak et al 2016). To evaluate fragment size, many surgeons now advocate obtaining a CT scan of the ankle in all patients with a known or suspected posterior malleolus fracture, although not yet part of British national guidelines (Solan & Sakellariou 2017). Recently, there has been a move towards fixation of more posterior malleolar fractures, as even with small areas of the articular surface there can be significant disruption to the posterior components of the syndesmosis (Bartoníček et al 2017). NICE emphasises the importance of early fixation by advising surgery on the day of injury or the day after in patients under 60 years of age (NICE 2016a).

Choice of surgery


Surgical treatment mostly takes the form of ORIF using plates and screws to reduce and stabilise the mortise (BOA 2016). Intraoperative fluoroscopy is used to monitor reduction and fixation (BOA 2016). Lateral malleolar fractures are most commonly fixed using an interfragmentary lag screw and a neutralisation plate (Lampridis et al 2018). However, if the fracture pattern allows more than one lag screw, a neutralisation plate is not mandatory (Lampridis et al 2018). Comminuted high-energy fibula fractures require stronger fixation with locking or reconstruction plates (Lampridis et al 2018). Alternatively, unstable distal fibular fractures can be treated using intramedullary fixation. This is a minimally invasive technique that has been shown to produce results comparable with plating, including a mean rate of union of 98.5% (Jain et al 2014).


Medial malleolar fractures, by comparison, are commonly fixed using lag screws or a tension band wire if the fracture fragment is small (Lampridis et al 2018). For comminuted medial

malleolar fractures, smaller sized screws are recommended. Another method used to treat vertical medial malleolar fractures specifically, involves the application of a buttress plate, or lag screws in combination with a plate (Lampridis et al 2018). Traditionally, posterior malleolar fractures have been stabilised using anteroposterior screw fixation, although many posterior malleolar fractures are managed nonoperatively in conjunction with surgically treated lateral or medial fractures (Odak et al 2016). Stabilising lateral or medial fractures inherently improves posterior fragment stability in these patients. More recently, there has been a trend towards fixation of posterior malleolus fractures using postero-lateral and postero-medial approaches which allow for more biomechanically sound fixation of the posterior malleolus (Gandham et al 2020).

After the ankle mortise is fixed, the syndesmosis must be assessed intraoperatively using a stress test such as the Hook test. This involves applying a lateral force to the distal fibula in order to obtain a radiograph of the mortise under stress (Stoffel et al 2009). If instability is identified, syndesmotic stabilisation is required. A number of options exist for stabilising the syndesmosis including screw fixation, with variation in the number of screws used and cortices crossed, as well as the use of TightRope® system. This system offers an alternative to conventional screw fixation that doesn't require removal and allows some movement at the syndesmosis. However, there is no clear evidence that one method is superior to another (Lampridis et al 2018).

Figure 3: Open reduction and internal fixation





Postoperative considerations

Physical and psychosocial complications

After surgery, patients are followed up within 6 weeks to assess the stability and alignment of the ankle and to monitor for any surgical complications (BOA 2016). It is especially important to be vigilant for postoperative complications amongst the elderly due to a high incidence of co-morbidities and poor bone quality (Srinivasan & Moran 2001).

Complications may include deep vein thrombosis and pulmonary embolism secondary to postoperative immobilisation and casting (Mehta et al 2014). Thromboprophylaxis should therefore be initiated in accordance with local guidelines and early mobilisation should be considered where appropriate as discussed below (BOA 2016).

Deep postoperative infection is a potentially limb-threatening complication that occurs in 3.4% of ankle fractures (Macera et al 2018). A high index of suspicion is, therefore, necessarily in order to facilitate early recognition and treatment. Management includes broad-spectrum IV antibiotics and debridement surgery (Zalavras et al 2009). Diabetic patients need careful perioperative monitoring as they are particularly prone to infection. A recent study reported that the rate of infection in operatively-treated diabetic ankle fracture patients varied from 11.5 to 21.5%, with higher rates in type 1 compared with type 2 diabetes and in open compared with closed fractures (Haddix et al 2018). Diabetic patients are also at an increased risk of other complications including impaired wound healing, malunion, non-union, Charcot arthropathy and soft tissue complications (Chaudhary et al 2008). To minimise these complications, meticulous perioperative glycaemic control with an insulin sliding scale is essential, as well as gentle soft tissue handling and robust fixation (Mehta et al 2014).

Later complications include malunion, non-union, secondary displacement and metalwork failure (Mehta et al 2014). Furthermore, patients are at risk of developing chronic pain, muscle atrophy and a stiff/swollen joint (Donken et al 2012). A long-term follow-up study reported that 63% of surgically-treated ankle fracture patients had stiffness, 45% had ankle

swelling, 50% experienced pain, and 38% had not returned to previous sporting levels, even 5 years after injury (Shah et al 2007). One of the most common long-term complications is post-traumatic osteoarthritis (PTOA). Advanced PTOA has been reported to affect 36.3% of surgically-treated malleolar fracture patients, at a mean of 18 years follow-up (Lübbecke et al 2012). It is reported that a lateral talar displacement of 1mm can reduce the contact area between talus and tibia by approximately 42%, which increases peak forces and cartilage wear and tear (Ramsey & Hamilton 1976). As surgical treatment of displaced fractures is more likely to restore normal anatomy, one may expect it to result in lower rates of PTOA. However, a Cochrane review comparing surgical and conservative treatment for ankle fractures concluded that there is insufficient evidence to definitively say which approach produces superior long-term outcomes (Donken et al 2012).

Ankle fractures may also have a significant psychosocial impact. A recent study assessed the experiences of 10 ankle fracture patients using semi-structured interviews between 19 weeks and 23 weeks following injury (McKeown et al 2020). These patients complained of a loss of independence, difficulties with activities of daily living, sleep disturbances, fatigue, depression and anxiety. These were attributable to factors such as reduced mobility, pain, skin issues and loss of strength and muscle bulk (McKeown et al 2020). Although this study included only a small number of patients and reported qualitative outcomes at a single point in time, it demonstrates the need for perioperative practitioners to take into account psychosocial factors when treating these individuals. Indeed, patients with ankle fractures may require psychological rehabilitation and social support in order to recover as quickly as possible (Mittly et al 2016). An improved understanding of these psychosocial factors will allow development of patient-tailored management plans.

Weight-bearing after surgery

Traditionally, the postoperative advice after ORIF has been to remain non-weight bearing for 6-8 weeks (King et al 2020). However, more recent evidence suggests that early mobilisation before this time period may accelerate recovery (Smeeing et al 2015). A number of multi-centred randomised controlled trials have studied this issue. In a study of 115 adult patients, the unprotected weight-bearing group had a significantly higher ankle function score than the protected weight-bearing and non-weight bearing groups at 6 weeks (Smeeing et al 2018). Moreover, the unprotected weight-bearing group had a significantly earlier return to work and sports with no difference in the complication rates. Lorente et al (2019) reported similar

1
2
3 272 findings in 70 elderly patients over a longer follow-up period. Specifically, this group found
4
5 273 that early weight-bearing patients scored higher than non-weight bearing patients using two
6
7 274 standardised measures of quality of life (SF-12 and Barthel Index), at 6-8 weeks, 1 year and 2
8
9 275 years of follow-up (Lorente et al 2019).

10 276
11
12 277 There is therefore an emerging evidence base in favour of weight-bearing after ankle fracture.
13
14 278 Indeed, current guidelines recommend that patients bear weight as tolerated in a splint/cast
15
16 279 post-surgery, except where the stability of fixation is uncertain or there are co-morbidities
17
18 280 such as peripheral neuropathy (BOA 2016). NICE also advises immediate, unrestricted
19
20 281 weight-bearing as tolerated for patients with stable fractures (NICE 2016a).

21 282
22 283 **Virtual fracture clinics**
23
24 284 Virtual Fracture Clinics (VFCs) are amongst the latest recommendations for research
25
26 285 published by NICE (NICE 2016a). According to the BOAST guidelines, “following acute
27
28 286 traumatic orthopaedic injury, patients should be seen in a new fracture clinic within 72 hours
29
30 287 of presentation” (BOA 2013). VFCs may help to meet this goal in the context of an
31
32 288 overstretched, and financially-limited traditional fracture clinic model (McKirdy &
33
34 289 Imbuldeniya 2017). The use of VFCs has been shown to significantly decrease non-attendees
35
36 290 and the number of days to first orthopaedic review, as well as significantly increase the
37
38 291 number of patients reviewed within 72 hours (McKirdy & Imbuldeniya 2017). Furthermore,
39
40 292 VFCs saved a local Clinical Commissioning Group £67,385.67 in their first year of use,
41
42 293 demonstrating clinical and cost-effectiveness (McKirdy & Imbuldeniya 2017). One concern
43
44 294 though, is that VFCs compromise the patient-doctor relationship. This has recently been
45
46 295 challenged by high patient satisfaction rates, with 94% of patients rating the service as good
47
48 296 or excellent, and 97% saying they were likely or extremely likely to recommend it to others
49
50 297 (Hawarden et al 2018)

51 298
52 299 **Conclusions**
53
54 300 Ankle fractures are complex orthopaedic injuries associated with significant morbidity and
55
56 301 mortality, especially in the perioperative period. It is therefore important to be conscious of
57
58 302 the key perioperative factors that influence the care of these patients.

59 303
60 304 **Declaration of conflicting interests**
305 The Authors declare no conflict of interest.

306

Funding

This research received no grant from any funding agency in the public, commercial or not-for-profit sectors.

310

References

Bachmann LM, Kolb E, Koller MT, Steurer J, Ter Riet G 2003 Accuracy of Ottawa ankle rules to exclude fractures of the ankle and mid-foot: Systematic review *Br Med J* 326(7386) 417–419. doi: 10.1136/bmj.326.7386.417.

315

Bartoniček J, Rammelt S, Tuček M 2017 Posterior Malleolar Fractures: Changing Concepts and Recent Developments *Foot Ankle Clin* 22(1) 125–145. doi: 10.1016/j.fcl.2016.09.009

318

BOA 2013 Fracture Clinic Services Available at

[https://www.boa.ac.uk/uploads/assets/7ded8f00-987e-42d5-a389e739b1e03b47/ec9d4564-4fa7-4d08-](https://www.boa.ac.uk/uploads/assets/7ded8f00-987e-42d5-a389e739b1e03b47/ec9d4564-4fa7-4d08-aef4efc3cede7d53/fracture%20clinic%20services.pdf?fbclid=IwAR3DGE2YDtl6nmPDKwwTwu6RTqH75DbZOHluOczjSijcF3gIMJ4u9khdZY)

[aef4efc3cede7d53/fracture%20clinic%20services.pdf?fbclid=IwAR3DGE2YDtl6nmPDKwwTwu6RTqH75DbZOHluOczjSijcF3gIMJ4u9khdZY](https://www.boa.ac.uk/uploads/assets/7ded8f00-987e-42d5-a389e739b1e03b47/ec9d4564-4fa7-4d08-aef4efc3cede7d53/fracture%20clinic%20services.pdf?fbclid=IwAR3DGE2YDtl6nmPDKwwTwu6RTqH75DbZOHluOczjSijcF3gIMJ4u9khdZY) [accessed March 2020]

324

BOA 2016 The Management of Ankle Fractures Available at

<https://www.boa.ac.uk/resources/boast-12-pdf.html> [accessed March 2020].

327

BOA & BAPRAS 2017 Open Fractures Available at

[https://www.boa.ac.uk/uploads/assets/3b91ad0a-9081-4253-](https://www.boa.ac.uk/uploads/assets/3b91ad0a-9081-4253-92f7d90e8df0fb2c/29bf80f11cb6-46b7-afc761119341447f/open%20fractures.pdf)

[92f7d90e8df0fb2c/29bf80f11cb6-46b7-afc761119341447f/open%20fractures.pdf](https://www.boa.ac.uk/uploads/assets/3b91ad0a-9081-4253-92f7d90e8df0fb2c/29bf80f11cb6-46b7-afc761119341447f/open%20fractures.pdf) [accessed March 2020].

332

Briet JP, Hietbrink F, Smeeing DP, Dijkgraaf MGW, Verleisdonk EJ, Houwert RM 2019

Ankle Fracture Classification: An Innovative System for Describing Ankle Fractures *J Foot Ankle Surg* 58(3) 492–496. doi: 10.1053/j.jfas.2018.09.028.

336

Bugler KE, Clement ND, Duckworth AD, White TO, McQueen MM, Court-Brown CM 2015

Open ankle fractures: who gets them and why? *Arch Orthop Trauma Surg* 135(3) 297–303.

doi: 10.1007/s00402-014-2140-3.

1
2
3 340
4
5 341 Chaudhary SB, Liporace FA, Gandi A, Donley BG, Pinzur MS, Lin SS 2008 Complications
6 342 of ankle fracture in patients with diabetes *J Am Acad Orthop Surg* 16(3) 159-170. doi:
7 343 10.5435/00124635-200803000-00007.
8
9 344
10
11 345 DeAngelis NA, Eskander MS, French BG 2007 Does medial tenderness predict deep deltoid
12 346 ligament incompetence in supination-external rotation type ankle fractures? *J Orthop*
13 347 *Trauma* 21(4) 244–247. doi: 10.1097/BOT.0b013e3180413835.
14
15 348
16
17 349 Donken CC, Al-Khateeb H, Verhofstad MH, Van Laarhoven CJ 2012 Surgical versus
18 350 conservative interventions for treating ankle fractures in adults *Cochrane Database Syst Rev*
19 351 2012(8) CD008470. doi: 10.1002/14651858.CD008470.pub2.
20
21 352
22
23 353 Elsoe R, Ostgaard SE, Larsen P 2018 Population-based epidemiology of 9767 ankle fractures
24 354 *Foot Ankle Surg* 24(1) 34–39. doi: 10.1016/j.fas.2016.11.002.
25
26 355
27
28 356 Gandham S, Millward G, Molloy A, Mason L 2020 Posterior malleolar fractures: A CT
29 357 guided incision analysis *The Foot* 43(), 101662. doi: 10.1016/j.foot.2019.101662
30 358
31 359 Haddix KP, Clement RC 3rd, Tennant JN, Ostrum RF 2018 Complications Following
32 360 Operatively Treated Ankle Fractures in Insulin- And Non-Insulin-Dependent Diabetic
33 361 Patients *Foot Ankle Spec* 11(3) 206-216.
34 362
35
36 363 Hsu RY, Lee Y, Hayda R, DiGiovanni CW, Mor V and Bariteau JT 2015 Morbidity and
37 364 mortality associated with geriatric ankle fractures *J Bone Joint Surg Am* 97(21) 1748–1755.
38 365 doi: 10.2106/JBJS.O.00095.
39 366
40
41 367 Jain S, Haughton BA, Brew C 2014 Intramedullary fixation of distal fibular fractures: a
42 368 systematic review of clinical and functional outcomes *J Orthop Traumatol* 15(4) 245-254.
43 369 doi: 10.1007/s10195-014-0320-0.
44
45 370
46
47 371 Jennison T, Brinsden M 2019 Fracture admission trends in England over a ten-year periods
48 372 *Ann R Coll Surg Eng* 101(3) 208-214. doi: 10.1308/rcsann.2019.0002.
49
50 373

- King CM, Doyle MD, Castellucci-Garza FM, Nguyentat A, Collman DR, Schuberth JM 2020 Early Protected Weightbearing After Open Reduction Internal Fixation of Ankle Fractures With Trans-syndesmotic Screws *J Foot Ankle Surg* S1067-2516(20) 30004-1. doi: 10.1053/j.jfas.2020.01.003.
- Lampridis V, Gougoulas N, Sakellariou A 2018 Stability in ankle fractures: Diagnosis and treatment *EFORT Open Rev* 3(5) 294–303. doi: 10.1302/2058-5241.3.170057.
- Lauge-hansen N 1954 Fractures of the ankle. III. Genetic roentgenologic diagnosis of fractures of the ankle *Am J Roentgenol Radium Ther Nucl Med* 71(3) 456–71.
- Leung KH, Fang CXS, Lau TW, Leung FKL 2016 Preoperative radiography versus computed tomography for surgical planning for ankle fractures *J Orthop Surg (Hong Kong)* 24(2) 158–162. doi: 10.1177/1602400207.
- Lorente A, Palacios P, Lorente R, Mariscal G, Barrios C, Gandía A 2020 Orthopedic treatment and early weight-bearing for bimalleolar ankle fractures in elderly patients: Quality of life and complications *Injury* 51(2) 548–553. doi: 10.1016/j.injury.2019.11.028.
- Lübbecke A, Salvo D, Stern R, Hoffmeyer P, Holzer N, Assal M 2012 Risk Factors for Post-Traumatic Osteoarthritis of the Ankle: An Eighteen Year Follow-up Study *Int Orthop* 36(7) 1403-1410. doi: 10.1007/s00264-011-1472-7.
- Macera A, Christian C, Sirleo L, Innocenti M 2018 Postoperative complications and reoperation rates following open reduction and internal fixation of ankle fracture *Joints* 6(2) 110–115. doi: 10.1055/s-0038-1653949.
- McKeown R, Kearney RS, Liew ZH, Ellard DR 2020 Patient experiences of an ankle fracture and the most important factors in their recovery: A qualitative interview study *BMJ Open* 10(2) e033539. doi: 10.1136/bmjopen-2019-033539.
- Medina McKeon JM, Hoch MC 2019 The ankle-joint complex: A kinesiological approach to lateral ankle sprains *J Athl Train* 54(6) 589–602. doi: 10.4085/1062-6050-472-17.


1
2
3 408 Mehta S, Rees K, Cutler L, Mangwani J 2014 Understanding risks and complications in the
4 409 management of ankle fractures *Indian J Orthop* 48(5) 445–452. doi: 10.4103/0019-
5 5413.139829.
6 410
7 411
8
9 412 Mittly V, Németh Z, Berényi K, Mintál T 2016 Mind Does Matter: The Psychological Effect
10 413 of Ankle Injury in Sport *J Psychol Psychother* 6(4) 1-7. doi: 10.4172/2161-0487.1000278.
11 414
12 415 Mordecai S, Al-Hadithy N 2011 Management of ankle fractures *BMJ* 343 d5204. doi:
13 416 10.1136/bmj.d5204.
14 417
15 418 Moore KL, Dalley AF, Agur AMR 2014 Clinically Oriented Anatomy 7th Edition Lippincott
16 419 Williams & Wilkins. doi: 10.1002/ca.22316.
17 420
18 421 NICE 2016a Fractures (non-complex): assessment and management Available at
19 422 <https://www.nice.org.uk/guidance/ng38> [accessed March 2020].
20 423
21 424 NICE 2016b Fractures (complex): assessment and management Available at
22 425 <https://www.nice.org.uk/guidance/ng37/chapter/Recommendations> [accessed March 2020].
23 426
24 427 Odak S, Ahluwalia R, Unnikrishnan P, Hennessy M, Platt S 2016 Management of Posterior
25 428 Malleolar Fractures: A Systematic Review *J Foot Ankle Surg* 55(1) 140-145. doi:
26 429 10.1053/j.jfas.2015.04.001.
27 430
28 431 Rammelt S, Endres T, Grass R, Zwipp H 2004 The role of external fixation in acute ankle
29 432 trauma *Foot Ankle Clin* 9(3) 455–474. doi: 10.1016/j.fcl.2004.05.001.
30 433
31 434 Ramsey PL, Hamilton W 1976 Changes in tibiotalar area of contact caused by lateral talar
32 435 shift *J Bone Joint Surg Am* 58(3) 356–357. doi: 10.2106/00004623-197658030-00010.
33 436
34 437 Sawant Y, Sanghvi D 2018 Magnetic resonance imaging of ankle ligaments: A pictorial
35 438 essay *Indian J Radiol Imaging* 28(4) 419–426. doi: 10.4103/ijri.IJRI_77_16.
36 439
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 440 Shah KN, Johnson JP, O'Donnell SW, Gil JA, Born CT, Hayda RA 2019 External Fixation
441 in the Emergency Department for Pilon and Unstable Ankle Fractures *J Am Acad Orthop*
442 *Surg* 27(12) e577-e584. doi: 10.5435/JAAOS-D-18-00080.
- 443
- 444 Shah NH, Sundaram RO, Velusamy A, Braithwaite IJ 2007 Five-year functional outcome
445 analysis of ankle fracture fixation *Injury* 38(11) 1308–1312. doi:
446 10.1016/j.injury.2007.06.002.
- 447
- 448 Slimmon D, Brukner P 2010 Sports ankle injuries - assessment and management *Aust Fam*
449 *Physician* 39(1–2) 18–22.
- 450
- 451 Smeeing DPJ, Houwert RM, Briet JP et al 2015 Weight-Bearing and Mobilization in the
452 Postoperative Care of Ankle Fractures: A Systematic Review and Meta-Analysis of
453 Randomized Controlled Trials and Cohort Studies *PLOS ONE* 10(2) e0118320. doi:
454 10.1371/journal.pone.0118320.
- 455
- 456 Smeeing DPJ, Houwert RM, Briet JP et al 2020 Weight-bearing or non-weight-bearing after
457 surgical treatment of ankle fractures: a multicenter randomized controlled trial *Eur J Trauma*
458 *Emerg S* 46(1) 121–130. doi: 10.1007/s00068-018-1016-6.
- 459
- 460 Solan M, Sakellariou A 2017. Posterior malleolus fractures: worth fixing. *Bone Joint J* 99-
461 B(11), 1413-1419. doi: 10.1302/0301-620x.99b11.bjj-2017-1072
- 462
- 463 Srinivasan CMS, Moran CG 2001 Internal fixation of ankle fractures in the very elderly
464 *Injury* 32(7) 559–563. doi: 10.1016/S0020-1383(01)00034-1.
- 465
- 466 Stiell IG, Greenberg GH, McKnight RD, Nair RC, McDowell I, Worthington JR 1992 A
467 study to develop clinical decision rules for the use of radiography in acute ankle injuries *Ann*
468 *Emerg Med* 21(4) 384–390. doi: 10.1016/S0196-0644(05)82656-3.
- 469
- 470 Stoffel K, Wysocki D, Baddour E, Nicholls R, Yates P 2009 Comparison of two
471 intraoperative assessment methods for injuries to the ankle syndesmosis. A cadaveric study. *J*
472 *Bone Joint Surg Am* 91(11) 2646-2652. doi: 10.2106/JBJS.G.01537.
- 473


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

474 Vangsness CT, Carter V, Newton E, Hunt T, Kerr R 1994 Radiographic Diagnosis of Ankle
475 Fractures: Are Three Views Necessary? *Foot Ankle Int* 15(4) 172–174. doi:
476 10.1177/107110079401500403.
477
478 Zalavras CG, Christensen T, Rigopoulos N, Holtom P, Patzakis MJ 2009 Infection following
479 operative treatment of ankle fractures *Clin Orthop Relat Res* 467(7) 1715–1720. doi:
480 10.1007/s11999-009-0743-8.

For Peer Review




Right foot in various views. (A) Lateral. (B) Medial. (C) Anterior. (D) Posterior (a part of the Achilles tendon has been removed). The names of the corresponding ligaments are given in blue.



Right foot viewed posteriorly. (A) Type A. (B) Type B. (C) Type C with disrupted syndesmosis.

Figure 3: Open reduction and internal fixation



Mortise view of an unstable lateral and medial malleolar ankle fracture before surgery (A) and after ORIF (B).

Mortise view of an unstable lateral and medial malleolar ankle fracture before surgery (A) and after ORIF (B).